arXiv:nlin/0408060v1 [nlin.SI] 31 Aug 2004

Toda equation and special polynomials associated with the Garnier system

Teruhisa TSUDA
Department of Mathematics, Kobe University,
Rokko, Kobe 657-8501, Japan.
e-mail: tudateru@ms.u-tokyo.ac.jp

Abstract

We prove that a certain sequence of tau functions of the Garnier system satisfies Toda equation. We construct a class of algebraic solutions of the system by the use of Toda equation; then show that the associated tau functions are expressed in terms of the universal character, which is a generalization of Schur polynomial attached to a pair of partitions.

This article is based on the results in the author's Ph.D thesis [19].

Introduction

The *Garnier system* is the following completely integrable Hamiltonian system of partial differential equations (see [1, 2, 4]):

$$\frac{\partial q_i}{\partial s_j} = \frac{\partial H_j}{\partial p_i}, \quad \frac{\partial p_i}{\partial s_j} = -\frac{\partial H_j}{\partial q_i}, \quad (i, j = 1, \ldots, N), \tag{0.1a}$$

with Hamiltonians

$$\begin{aligned} s_i(s_i-1)H_i &= q_i\left(\alpha + \sum_j q_jp_j\right)\left(\alpha + \kappa_\infty + \sum_j q_jp_j\right) + s_ip_i(q_ip_i - \theta_i) \\ &\quad - \sum_{j(\neq i)} R_{ji}(q_jp_j - \theta_j)q_ip_j - \sum_{j(\neq i)} S_{ij}(q_ip_i - \theta_i)q_jp_i \\ &\quad - \sum_{j(\neq i)} R_{ij}q_jp_j(q_ip_i - \theta_i) - \sum_{j(\neq i)} R_{ij}q_ip_i(q_jp_j - \theta_j) \\ &\quad -(s_i+1)(q_ip_i - \theta_i)q_ip_i + (\kappa_1 s_i + \kappa_0 - 1)q_ip_i, \end{aligned} \tag{0.1b}$$

where $R_{ij} = s_i(s_j - 1)/(s_j - s_i)$, $S_{ij} = s_i(s_i - 1)/(s_i - s_j)$ and

$$\alpha = -\frac{1}{2}\left(\kappa_0 + \kappa_1 + \kappa_\infty + \sum_i \theta_i - 1\right). \tag{0.2}$$

Here the symbols $\sum_i$ and $\sum_{i(\neq j)}$ stand for the summation over $i = 1, \ldots, N$ and over $i = 1, \ldots, j-1, j+1, \ldots, N$, respectively. System (0.1) contains $N + 3$ constant parameters

$$\vec{\kappa} = (\kappa_0, \kappa_1, \kappa_\infty, \theta_1, \ldots, \theta_N) \in \mathbb{C}^{N+3}, \tag{0.3}$$

so that we often denote it by $\mathcal{H}_N = \mathcal{H}_N(\vec{\kappa}) = \mathcal{H}_N(q, p, s, H; \vec{\kappa})$, and so on. The Garnier system governs the monodromy preserving deformation of a Fuchsian differential equation with $N + 3$ singularities and is an extension of the sixth Painlevé equation P_{VI}; for $N = 1$, (0.1) is equivalent to the Hamiltonian system of P_{VI} (see [13]), in fact.

In this paper, we prove that a certain sequence of τ-functions of the Garnier system satisfies Toda equation. We construct a class of algebraic solutions of the system by using Toda equation; then show that the corresponding τ-functions are expressed in terms of the universal character, which is a generalization of Schur polynomial attached to a pair of partitions.

First we introduce a group of birational canonical transformations of the Garnier system $\mathcal{H}_N$. The group forms an infinite group which contains a translation $\mathbb{Z}$; see Sect. 1. We define a function $\tau = \tau(s; \vec{\kappa})$, called the *$\tau$-function* (see [2, 4]), by

$$\mathrm{d}\log\tau = \sum_i H_i \mathrm{d}s_i. \tag{0.4}$$

By the use of birational symmetries of $\mathcal{H}_N$, we have the

Theorem 0.1. *A certain sequence of τ-functions $\{\tau_n | n \in \mathbb{Z}\}$ satisfies the Toda equation:*

$$XY \log \tau_n = c_n \frac{\tau_{n-1}\tau_{n+1}}{\tau_n^2}, \tag{0.5}$$

where X, Y being vector fields such that $[X, Y] = 0$ and c_n a nonzero constant.

(See Theorem 2.2.)

Consider the fixed point of a certain birational symmetry, we obtain an algebraic solution of the Garnier system. For example, if $\kappa_0 = \kappa_1 = 1/2$, then $\mathcal{H}_N$ admits an algebraic solution

$$(q_i, p_i) = \left(\frac{\theta_i \sqrt{s_i}}{\kappa_\infty}, \frac{\kappa_\infty}{2\sqrt{s_i}} \right), \quad i = 1, \dots, N. \tag{0.6}$$

Applying the action of the group of birational symmetries, we thus have the

Theorem 0.2. *If two components of the parameter $\vec{\kappa} = (\kappa_0, \kappa_1, \kappa_\infty, \theta_1, \dots, \theta_N)$ are half integers then the Garnier system $\mathcal{H}_N$ admits an algebraic solution.*

(See Theorem 3.1.)

Secondly we investigate the τ-functions associated with algebraic solutions of the Garnier system. Starting from the τ-function corresponding to an algebraic solution, we determine a sequence of τ-functions by means of Toda equation. Such a sequence of τ-functions is converted to polynomials $T_{m,n} = T_{m,n}(t)$ $(m, n \in \mathbb{Z})$ through a certain normalization, where $t = (t_1, \dots, t_N)$ and $t_i = \sqrt{s_i}$. We call $T_{m,n}$ *special polynomials* associated with algebraic solutions of $\mathcal{H}_N$ (see Sect. 3). Algebraic solutions are explicitly written in terms of the special polynomials.

Theorem 0.3. *If $\kappa_0 = 1/2 + m + n$, $\kappa_1 = 1/2 + m - n$ $(m, n \in \mathbb{Z})$, then $\mathcal{H}_N$ admits an algebraic solution given by*

$$\begin{aligned} q_i &= \frac{t_i \dfrac{\partial}{\partial t_i} \log \dfrac{T_{m+1,n}}{T_{m,n+1}}}{\displaystyle\sum_j t_j \frac{\partial}{\partial t_j} \log \frac{T_{m+1,n}}{T_{m,n+1}} - 2m + 2n - 1}, \\ 2q_i p_i &= \theta_i + m + n + t_i \frac{\partial}{\partial t_i} \log \frac{T_{m,n}}{T_{m,n+1}}. \end{aligned} \tag{0.7}$$

(See Theorem 3.3.) Note that we immediately obtain also the expressions of the other algebraic solutions in Theorem 0.2, via the birational symmetries of $\mathcal{H}_N$. Finally we give an explicit formula for $T_{m,n}$ in terms of the universal character (see [7, 16]), which is a generalization of Schur polynomial.

Theorem 0.4. *The special polynomials $T_{m,n}(t)$ $(m, n \in \mathbb{Z})$ is expressed as follows:*

$$T_{m,n}(t) = N_{m,n} S_{[\lambda,\mu]}(x, y). \tag{0.8}$$

Here $S_{[\lambda,\mu]}(x,y) = S_{[\lambda,\mu]}(x_1, x_2, \dots, y_1, y_2, \dots)$ *denotes the universal character attached to a pair of partitions*

$$\lambda = (u, u-1, \dots, 2, 1), \quad \mu = (v, v-1, \dots, 2, 1), \tag{0.9}$$

with $u = |n-m-1/2| - 1/2$, $v = |n+m-1/2| - 1/2$; $N_{m,n}$ *is a certain normalization factor, and*

$$x_n = \frac{-\kappa_\infty + \sum_i \theta_i t_i^n}{n}, \quad y_n = \frac{-\kappa_\infty + \sum_i \theta_i t_i^{-n}}{n}. \tag{0.10}$$

(See Theorem 3.5 and also Corollary 3.6.) Recall that the universal character is the irreducible character of a rational representation of $GL(n)$, while Schur polynomial that of a polynomial representation; see [7]. Hence Theorem 0.4 shows us a relationship between the representation theory of $GL(n)$ and the Garnier system, or the theory of monodromy preserving deformation.

We propose in [16] an infinite dimensional integrable system characterized by the universal characters, called the UC hierarchy; and regard it as an extension of the KP hierarchy. Since all the universal characters are solutions of the UC hierarchy, it would be an interesting problem to construct a certain reduction procedure from the hierarchy to the Garnier system; *cf.* [18].

In Sect. 1, we present a group of birational canonical transformations of the Garnier system $\mathcal{H}_N$. In Sect. 2, we prove that a certain sequence of τ-functions satisfies Toda equation. In Sect. 3, we construct a class of algebraic solutions of $\mathcal{H}_N$ by using Toda equation; then show that the associated τ-functions are explicitly written in terms of the universal characters. Sect. 4 is devoted to the verification of Theorem 3.5.

1 Birational symmetry

First we introduce a group of birational canonical transformations of the Garnier system $\mathcal{H}_N(\vec{\kappa})$; then see that it forms an infinite group which contains a translation $\mathbb{Z}$.

It is known that $\mathcal{H}_N$ has a symmetry which is isomorphic to the symmetric group.

Theorem 1.1 (see [2, 5]**).** *The Garnier system* $\mathcal{H}_N(\vec{\kappa})$ *has birational canonical transformations*

$$\sigma_m : (q, p, s, \vec{\kappa}) \mapsto (Q, P, S, \sigma_m(\vec{\kappa})), \quad 1 \le m \le N+2,$$

given in the following table:

σ_m	action on $\vec{\kappa}$	Q_i	P_i	S_i
σ_m $(m \leq N)$	$\theta_m \leftrightarrow \kappa_0$	$Q_i = \dfrac{q_i}{R_{im}} \quad (i \neq m),$ $Q_m = \dfrac{s_m(1-g_s)}{s_m - 1}$	$P_i = R_{im}\left(p_i - \dfrac{s_m}{s_i}p_m\right),$ $P_m = -(s_m - 1)p_m$	$S_i = \dfrac{s_m - s_i}{s_m - 1},$ $S_m = \dfrac{s_m}{s_m - 1}$
σ_{N+1}	$\kappa_1 \leftrightarrow \kappa_0$	$Q_i = \dfrac{q_i}{s_i}$	$P_i = s_i p_i$	$S_i = \dfrac{1}{s_i}$
σ_{N+2}	$\kappa_1 \leftrightarrow \kappa_\infty$	$Q_i = \dfrac{q_i}{g_1 - 1}$	$P_i = (g_1 - 1)$ $\times \left(p_i - \alpha - \sum_j q_j p_j\right)$	$S_i = \dfrac{s_i}{s_1 - 1}$

where $g_1 = \sum_j q_j$, $g_s = \sum_j q_j/s_j$, *and* $\langle \sigma_1, \ldots, \sigma_{N+2} \rangle \simeq \mathfrak{S}_{N+3}$.

Theorem 1.1 is verified by considering a permutation among $N+3$ singularities of the associated linear differential equation; see [2, 5]. Combine the above $\mathfrak{S}_{N+3}$-symmetry with the fact that Hamiltonians H_i (see (0.1b)) are invariant under the action

$$\kappa_\infty \mapsto -\kappa_\infty,$$

we obtain also the following birational transformations.

Theorem 1.2. *The Garnier system* $\mathcal{H}_N(\vec{\kappa})$ *has the birational canonical transformations*

$$R_\Delta : \mathcal{H}_N(\vec{\kappa}) \to \mathcal{H}_N(R_\Delta(\vec{\kappa})).$$

Here the birational transformations $R_\Delta : (q, p) \mapsto (Q, P)$ *are described as follows*:

R_Δ	action on $\vec{\kappa}$	Q_i	P_i
R_{κ_∞}	$\kappa_\infty \mapsto -\kappa_\infty$	$Q_i = q_i$	$P_i = p_i$
R_{κ_1}	$\kappa_1 \mapsto -\kappa_1$	$Q_i = q_i$	$P_i = p_i - \dfrac{\kappa_1}{g_1 - 1}$
R_{κ_0}	$\kappa_0 \mapsto -\kappa_0$	$Q_i = q_i$	$P_i = p_i - \dfrac{\kappa_0}{s_i(g_s - 1)}$
R_{θ_j}	$\theta_j \mapsto -\theta_j$	$Q_i = q_i$	$P_j = p_j - \frac{\theta_j}{q_j}, \quad P_i = p_i \quad (i \neq j)$

We now introduce another birational transformation of $\mathcal{H}_N(\vec{\kappa})$ which seems to be more nontrivial than the previous ones.

Theorem 1.3. *The Garnier system* $\mathcal{H}_N(\vec{\kappa})$ *has the birational canonical transformation*

$$R_\tau : \mathcal{H}_N(q, p, s, H; \vec{\kappa}) \to \mathcal{H}_N(Q, P, s, \widetilde{H}; R_\tau(\vec{\kappa})),$$

where $R_\tau(\vec{\kappa}) = (-\kappa_0+1, -\kappa_1+1, -\kappa_\infty, -\theta_1, \ldots, -\theta_N)$ *and*

$$\begin{aligned} Q_i &= \frac{s_i p_i(q_i p_i - \theta_i)}{\left(\alpha + \sum_j q_j p_j\right)\left(\alpha + \kappa_\infty + \sum_j q_j p_j\right)}, && \text{(1.1a)} \\ Q_i P_i &= -q_i p_i, && \text{(1.1b)} \\ \widetilde{H}_i &= H_i - \frac{q_i p_i}{s_i}. && \text{(1.1c)} \end{aligned}$$

Let G be a group of birational canonical transformations of $\mathcal{H}_N(\vec{\kappa})$ defined by

$$G = \langle \sigma_1, \ldots, \sigma_{N+2}, R_{\kappa_0}, R_{\kappa_1}, R_{\kappa_\infty}, R_{\theta_1}, \ldots, R_{\theta_N}, R_\tau \rangle. \tag{1.2}$$

We see that G forms an infinite group which contains $\mathbb{Z}$. For instance, let

$$l = R_{\kappa_1} \circ R_\tau \circ R_{\theta_1} \circ \cdots \circ R_{\theta_N} \circ R_{\kappa_\infty} \circ R_{\kappa_0} \in G,$$

then l acts on the parameter as its translation:

$$l(\vec{\kappa}) = \vec{\kappa} + (1, -1, 0, 0, \ldots, 0),$$

thus $\{l^n\} \simeq \mathbb{Z} \subset G$.

Remark 1.4. Group G might not fill all the birational symmetries of $\mathcal{H}_N$. If $\theta_i = 0$ $(i \neq 1)$, then $\mathcal{H}_N$ admits a particular solution written in terms of solutions of the sixth Painlevé equation P_{VI}; see [14, Theorem 6.1]. However group G with the restriction to $\theta_i = 0$ $(i \neq 1)$ does not form the affine Weyl group of type $D_4^{(1)}$, which is the group of birational symmetries for P_{VI} ; see [13]. So the author suspects that there would exist another hidden symmetry of $\mathcal{H}_N$. Anyway, it is an important problem to determine the group of all birational symmetries of the Garnier system $\mathcal{H}_N$.

Proof of Theorem 1.3. First we shall verify that the transformation R_τ is a canonical transformation of Hamiltonian system $\mathcal{H}_N$; that is,

$$\sum_i (\mathrm{d}p_i \wedge \mathrm{d}q_i - \mathrm{d}H_i \wedge \mathrm{d}s_i) = \sum_i \left(\mathrm{d}P_i \wedge \mathrm{d}Q_i - \mathrm{d}\widetilde{H}_i \wedge \mathrm{d}s_i\right). \tag{1.3}$$

From (1.1b), we have

$$P_i \mathrm{d}Q_i + Q_i \mathrm{d}P_i = -p_i \mathrm{d}q_i - q_i \mathrm{d}p_i. \tag{1.4}$$

Consider the logarithmic derivative of (1.1a), we have

$$\begin{aligned} \frac{\mathrm{d}Q_i}{Q_i} &= \frac{\mathrm{d}s_i}{s_i} + \frac{\mathrm{d}p_i}{p_i} + \frac{p_i \mathrm{d}q_i + q_i \mathrm{d}p_i}{q_i p_i - \theta_i} \\ &\quad - \left(\frac{1}{\alpha + \sum_j q_j p_j} + \frac{1}{\alpha + \kappa_\infty + \sum_j q_j p_j}\right) \sum_j \mathrm{d}(q_j p_j). \end{aligned} \tag{1.5}$$

By taking the wedge product of (1.4) and (1.5), we obtain

$$\begin{aligned}\mathrm{d}P_i \wedge \mathrm{d}Q_i &= \mathrm{d}p_i \wedge \mathrm{d}q_i - \mathrm{d}\left(\frac{q_i p_i}{s_i}\right) \wedge \mathrm{d}s_i \\ &\quad + \left(\frac{1}{\alpha + \sum_j q_j p_j} + \frac{1}{\alpha + \kappa_\infty + \sum_j q_j p_j}\right) \mathrm{d}(q_i p_i) \wedge \sum_{j(\neq i)} \mathrm{d}(q_j p_j);\end{aligned}$$

hence

$$\sum_i \mathrm{d}P_i \wedge \mathrm{d}Q_i = \sum_i \mathrm{d}p_i \wedge \mathrm{d}q_i - \sum_i \mathrm{d}\left(\frac{q_i p_i}{s_i}\right) \wedge \mathrm{d}s_i. \tag{1.6}$$

On the other hand, it follows from (1.1c) that

$$\mathrm{d}\widetilde{H}_i \wedge \mathrm{d}s_i = \mathrm{d}H_i \wedge \mathrm{d}s_i - \mathrm{d}\left(\frac{q_i p_i}{s_i}\right) \wedge \mathrm{d}s_i. \tag{1.7}$$

Combining (1.6) and (1.7), we get (1.3).

Secondly we shall prove that

$$\widetilde{H}_i = H_i(Q, P, s, R_\tau(\vec{\kappa})). \tag{1.8}$$

Notice that $s_j S_{ij} = s_i R_{ji}$. By using (1.1a) and (1.1b) we have the formulae:

$$Q_i\left(-\alpha + \sum_j Q_j P_j\right)\left(-\alpha - \kappa_\infty + \sum_j Q_j P_j\right) = s_i p_i (q_i p_i - \theta_i), \tag{1.9a}$$

$$s_i P_i (Q_i P_i + \theta_i) = q_i \left(\alpha + \sum_j q_j p_j\right)\left(\alpha + \kappa_\infty + \sum_j q_j p_j\right), \tag{1.9b}$$

$$\sum_{j(\neq i)} R_{ji}(Q_j P_j + \theta_j) Q_i P_j = \sum_{j(\neq i)} S_{ij}(q_i p_i - \theta_i) q_j p_i, \tag{1.9c}$$

$$\sum_{j(\neq i)} S_{ij}(Q_i P_i + \theta_i) Q_j P_i = \sum_{j(\neq i)} R_{ji}(q_j p_j - \theta_j) q_i p_j. \tag{1.9d}$$

Recall the definition of Hamiltonian H_i; see (0.1b). Then we verify (1.8) by (1.9) immediately. The proof is now complete. ■

2 Toda equation

In this section we show that a certain sequence of τ-functions satisfies the Toda equation.

Since the 1-form $\omega = \sum_i H_i \mathrm{d}s_i$ is closed, we can define, up to multiplicative constants, a function $\tau = \tau(s; \vec{\kappa})$ called the *τ-function* by (see [2, 4])

$$\mathrm{d}\log\tau = \sum_i H_i \mathrm{d}s_i. \tag{2.1}$$

Let l be a birational canonical transformation of $\mathcal{H}_N$ defined by

$$l = R_{\kappa_1} \circ R_\tau \circ R_{\theta_1} \circ \cdots \circ R_{\theta_N} \circ R_{\kappa_\infty} \circ R_{\kappa_0}, \tag{2.2}$$

then l acts on the parameter $\vec{\kappa} = (\kappa_0, \kappa_1, \kappa_\infty, \theta_1, \ldots, \theta_N)$ as its translation:

$$l(\vec{\kappa}) = \vec{\kappa} + (1, -1, 0, 0, \ldots, 0).$$

Let $(q_i(s), p_i(s), H_i(s))$ be a solution of the Garnier system $\mathcal{H}_N(\vec{\kappa})$ and set

$$\begin{aligned} (q_i^+, p_i^+, H_i^+) &= (l(q_i), l(p_i), l(H_i)), \\ (q_i^-, p_i^-, H_i^-) &= (l^{-1}(q_i), l^{-1}(p_i), l^{-1}(H_i)), \end{aligned} \tag{2.3}$$

then we have the

Proposition 2.1. *The triple of Hamiltonians* $(H_i^+(s), H_i(s), H_i^-(s))$ *satisfies the differential equation*:

$$H_i^+ - 2H_i + H_i^- = \frac{\partial}{\partial s_i} \log F(s), \tag{2.4}$$

where

$$F(s) = \left(\sum_j (s_j - 1) \frac{\partial}{\partial s_j} - 1 \right) \sum_k s_k(s_k - 1) H_k - \kappa_1(\kappa_0 - 1) + \alpha(\alpha + \kappa_\infty). \tag{2.5}$$

One can prove the proposition by straightforward computations, via the birational transformations given in Sect. 1; see [19], for details.

Let $\tau^\pm = l^{\pm 1}(\tau)$, then we rewrite (2.4) into

$$\left(\sum_i (s_i - 1) \frac{\partial}{\partial s_i} - 1 \right) \left(\sum_j s_j(s_j - 1) \frac{\partial}{\partial s_j} \right) \log \tau - \kappa_1(\kappa_0 - 1) + \alpha(\alpha + \kappa_\infty) = c \frac{\tau^+ \tau^-}{\tau^2}, \tag{2.6}$$

where c is a nonzero constant. Consider the change of variables $s_i = \xi_i/(\xi_i - 1)$ and the differential operators:

$$A = \sum_i \xi_i \frac{\partial}{\partial \xi_i}, \quad B = \sum_i \frac{\partial}{\partial \xi_i}, \tag{2.7}$$

then we have

$$\left(\sum_i (s_i - 1) \frac{\partial}{\partial s_i} - 1 \right) \left(\sum_j s_j(s_j - 1) \frac{\partial}{\partial s_j} \right) = (A - B + 1)A. \tag{2.8}$$

Note that

$$[A, B] = AB - BA = -B. \tag{2.9}$$

Let

$$\psi = \Delta^{\frac{2}{N(N-1)}}, \tag{2.10}$$

where Δ denotes the difference product of $(\xi_1, \xi_2, \ldots, \xi_N)$, *i.e.*,

$$\Delta = \prod_{i>j} (\xi_i - \xi_j) = \begin{vmatrix} 1 & 1 & \cdots & 1 \\ \xi_1 & \xi_2 & \cdots & \xi_N \\ \vdots & \vdots & \ddots & \vdots \\ \xi_1^{N-1} & \xi_2^{N-1} & \cdots & \xi_N^{N-1} \end{vmatrix}.$$

Since

$$A\Delta = \frac{N(N-1)}{2}\Delta, \quad B\Delta = 0,$$

we have

$$A\psi = \psi, \quad B\psi = 0. \tag{2.11}$$

Introduce the vector fields

$$X = \psi(A - B), \quad Y = \psi A. \tag{2.12}$$

One can easily verify that $[X, Y] = 0$,

$$XY = \psi^2 (A - B + 1)A, \tag{2.13}$$

and

$$XY \log \psi = \psi^2. \tag{2.14}$$

by using (2.9) and (2.11).

Let us consider the sequence of τ-functions $\{\tau_n | n \in \mathbb{Z}\}$ defined by

$$\tau_n = \psi^{a_n} l^n(\tau), \tag{2.15}$$

with

$$a_n = -(\kappa_1 - n)(\kappa_0 + n - 1) + \alpha(\alpha + \kappa_\infty). \tag{2.16}$$

Substitute (2.15) into (2.6), by virtue of (2.13) and (2.14), we now arrive at the

Theorem 2.2. *The sequence $\{\tau_n | n \in \mathbb{Z}\}$ satisfies the Toda equation*:

$$XY \log \tau_n = c_n \frac{\tau_{n-1}\tau_{n+1}}{\tau_n^2}, \tag{2.17}$$

where X, Y being vector fields such that $[X, Y] = 0$ and c_n a nonzero constant.

Remark 2.3. A sequence of τ-functions corresponding to other translations also satisfies the Toda equation. For instance, let us consider the birational transformation $\widetilde{l}$ defined by

$$\widetilde{l} = R_{\kappa_1} \circ l \circ R_{\kappa_1}, \tag{2.18}$$

which acts on the parameter $\vec{\kappa}$ as its translation:

$$\widetilde{l}(\vec{\kappa}) = \vec{\kappa} + (1, 1, 0, 0, \ldots, 0).$$

It is easy to see that

$$R_{\kappa_1}(\tau)=\tau\prod_i(s_i-1)^{-\kappa_1\theta_i}. \tag{2.19}$$

Combine this with (2.6), we obtain

$$\left(\sum_i\frac{\partial}{\partial s_i}-1\right)\left(\sum_j s_j(s_j-1)\frac{\partial}{\partial s_j}\right)\log\tau+\alpha(\alpha+\kappa_\infty)=c\frac{\widetilde{l}^{-1}(\tau)\,\widetilde{l}(\tau)}{\tau^2}. \tag{2.20}$$

Also (2.20) is equivalent to the Toda equation via a similar change of variables as above.

3 Algebraic solutions in terms of universal characters

In this section we construct a class of algebraic solutions of the Garnier system $\mathcal{H}_N$ and then express it in terms of the universal characters.

3.1 Algebraic solutions

Consider the birational canonical transformation

$$w_0=R_\tau\circ R_{\theta_1}\circ\cdots\circ R_{\theta_N}\circ R_{\kappa_\infty}, \tag{3.1}$$

given as follows:

$$w_0:\mathcal{H}_N(q,p;\vec{\kappa})\to\mathcal{H}_N(Q,P;w_0(\vec{\kappa})),$$

where $w_0(\vec{\kappa})=(-\kappa_0+1,-\kappa_1+1,\kappa_\infty,\theta_1,\dots,\theta_N)$ and

$$Q_i = \frac{s_ip_i(q_ip_i-\theta_i)}{\left(\alpha+\sum_j q_jp_j\right)\left(\alpha+\kappa_\infty+\sum_j q_jp_j\right)}, \tag{3.2a}$$

$$Q_iP_i = -q_ip_i+\theta_i. \tag{3.2b}$$

If $\kappa_0=\kappa_1=1/2$, the fixed point with respect to the action of w_0 is

$$(q_i,p_i)=\left(\frac{\theta_i\sqrt{s_i}}{\kappa_\infty},\frac{\kappa_\infty}{2\sqrt{s_i}}\right),\quad i=1,\dots,N. \tag{3.3}$$

This is an algebraic solution of $\mathcal{H}_N$. Applying the birational symmetries G (see Sect. 1) to (3.3), we obtain a class of algebraic solutions.

Theorem 3.1. *If two components of the parameter* $\vec{\kappa}=(\kappa_0,\kappa_1,\kappa_\infty,\theta_1,\dots,\theta_N)$ *are half integers then* $\mathcal{H}_N$ *admits an algebraic solution.*

3.2 Special polynomials

Substituting the algebraic solution, (3.3), into Hamiltonians (see (0.1b)), we have

$$s_i(s_i-1)H_i = -\frac{1}{2}\kappa_\infty\theta_i\sqrt{s_i} + \frac{1}{4}\theta_i(s_i-1) + \frac{1}{2}\sum_j \theta_i\theta_j\frac{\sqrt{s_is_j}+1}{\sqrt{s_j/s_i}+1}; \tag{3.4}$$

and then the corresponding τ-function is given as follows:

$$\tau_{0,0} = \prod_i s_i^{-\theta_i(\theta_i-1)/4}(\sqrt{s_i}+1)^{\theta_i(\sum_k\theta_k+\kappa_\infty)/2}(\sqrt{s_i}-1)^{\theta_i(\sum_k\theta_k-\kappa_\infty)/2}\prod_{i,j}(\sqrt{s_i}+\sqrt{s_j})^{-\theta_i\theta_j/2}. \tag{3.5}$$

Let us consider the birational transformations l and $\widetilde{l}$, defined respectively by (2.2) and (2.18), which act on the parameter $\vec{\kappa}$ as its translations:

$$\begin{aligned} l(\vec{\kappa}) &= \vec{\kappa} + (1,-1,0,0,\ldots,0), \\ \widetilde{l}(\vec{\kappa}) &= \vec{\kappa} + (1,1,0,0,\ldots,0). \end{aligned} \tag{3.6}$$

Introduce a family of τ-functions $\tau_{m,n}$ $(m,n\in\mathbb{Z})$ defined by

$$\widetilde{l}^m l^n(\tau_{0,0}) = \tau_{m,n}. \tag{3.7}$$

Let

$$s_i = t_i^2, \tag{3.8}$$

then (3.5) is rewritten as

$$\tau_{0,0} = \prod_i t_i^{-\theta_i(\theta_i-1)/2}(t_i+1)^{\theta_i(\sum_k\theta_k+\kappa_\infty)/2}(t_i-1)^{\theta_i(\sum_k\theta_k-\kappa_\infty)/2}\prod_{i,j}(t_i+t_j)^{-\theta_i\theta_j/2}. \tag{3.9a}$$

Applying the action of $\widetilde{l}$ and l, we see that

$$\begin{aligned} \tau_{0,1} &= \prod_i t_i^{-\theta_i}\tau_{0,0}, && (3.9\text{b}) \\ \tau_{1,0} &= \left(\prod_i t_i^{-\theta_i}(t_i+1)^{\theta_i}(t_i-1)^{\theta_i}\right)\left(\sum_j\theta_jt_j-\kappa_\infty\right)\tau_{0,0}, && (3.9\text{c}) \\ \tau_{1,1} &= \left(\prod_i t_i^{-2\theta_i}(t_i+1)^{\theta_i}(t_i-1)^{\theta_i}\right)\left(\kappa_\infty-\sum_j\theta_jt_j^{-1}\right)\tau_{0,0}. && (3.9\text{d}) \end{aligned}$$

The τ-functions, $\tau_{m,n}$ $(m,n\in\mathbb{Z})$, are determined successively by the use of the Toda equations (2.6) and (2.20), from the above initial values (3.9).

Now let us define the functions, $T_{m,n} = T_{m,n}(t)$ $(m,n\in\mathbb{Z})$, by

$$\begin{aligned} T_{m,n}(t) &= \tau_{m,n}\prod_i\Big\{t_i^{(\theta_i+m+n)(\theta_i+m+n-1)/2}(t_i+1)^{-\theta_i(\sum_k\theta_k+\kappa_\infty+2m)/2} \\ &\qquad\times(t_i-1)^{-\theta_i(\sum_k\theta_k-\kappa_\infty+2m)/2}\Big\}\prod_{i,j}(t_i+t_j)^{\theta_i\theta_j/2}. \end{aligned} \tag{3.10}$$

Substituting (3.10) into (2.6) and (2.20) with $c = 1/4$, we thus obtain the recurrence relations for $T_{m,n}$.

Proposition 3.2. *The function* $T_{m,n} = T_{m,n}(t)$ $(m, n \in \mathbb{Z})$ *satisfies the following recurrence relations*:

$$\begin{aligned} T_{m+1,n} &= \prod_i t_i \Bigg\{ \left(\sum_i \frac{t_i^2 - 1}{t_i} \frac{\partial}{\partial t_i} - 2 \right) \sum_i t_i (t_i^2 - 1) \frac{\partial}{\partial t_i} \log T_{m,n} \\ &\quad + \kappa_\infty \sum_i \theta_i \frac{t_i^2 + 1}{t_i} - \frac{1}{2} \sum_{i,j} \theta_i \theta_j \frac{t_i^2 + t_j^2}{t_i t_j} - \kappa_\infty^2 + (2m)^2 \Bigg\} \frac{{T_{m,n}}^2}{T_{m-1,n}}, \qquad (3.11a) \\ T_{m,n+1} &= \prod_i t_i \Bigg\{ \left(\sum_i \frac{t_i^2 - 1}{t_i} \frac{\partial}{\partial t_i} - 2 \right) \sum_i t_i (t_i^2 - 1) \frac{\partial}{\partial t_i} \log T_{m,n} \\ &\quad + \kappa_\infty \sum_i \theta_i \frac{t_i^2 + 1}{t_i} - \frac{1}{2} \sum_{i,j} \theta_i \theta_j \frac{t_i^2 + t_j^2}{t_i t_j} - \kappa_\infty^2 + (2n-1)^2 \Bigg\} \frac{{T_{m,n}}^2}{T_{m,n-1}}. \qquad (3.11b) \end{aligned}$$

Here the initial values are given as follows:

$$T_{0,0} = T_{0,1} = 1, \quad T_{1,0} = \sum_i \theta_i t_i - \kappa_\infty, \quad T_{1,1} = \prod_i t_i \left(\kappa_\infty - \sum_j \theta_j t_j^{-1} \right). \qquad (3.12)$$

We call $T_{m,n}(t)$ *special polynomials* associated with algebraic solutions of $\mathcal{H}_N$. By the above recurrence relations (3.11), we can only state that $T_{m,n}(t)$ are rational functions in $t = (t_1, \ldots, t_N)$. We will show that $T_{m,n}(t)$ are indeed polynomials; see Theorem 3.5 and Corollary 3.6 below. Note that

$$T_{-m,n}(t) = T_{m,1-n}(t) = (-1)^{m(2n-1)} \prod_i t_i^{m^2 + n(n-1)} T_{m,n}(t^{-1}), \qquad (3.13)$$

which is verified easily by the recurrence relations and initial values. Algebraic solutions of $\mathcal{H}_N$ are explicitly written in terms of the special polynomials $T_{m,n}(t)$.

Theorem 3.3. *If* $\kappa_0 = 1/2 + m + n$, $\kappa_1 = 1/2 + m - n$ $(m, n \in \mathbb{Z})$, *then* $\mathcal{H}_N$ *admits an algebraic solution given as follows*:

$$\begin{aligned} q_i &= \frac{t_i \dfrac{\partial}{\partial t_i} \log \dfrac{T_{m+1,n}}{T_{m,n+1}}}{\displaystyle\sum_j t_j \frac{\partial}{\partial t_j} \log \frac{T_{m+1,n}}{T_{m,n+1}} - 2m + 2n - 1}, \qquad (3.14a) \\ 2 q_i p_i &= \theta_i + m + n + t_i \frac{\partial}{\partial t_i} \log \frac{T_{m,n}}{T_{m,n+1}}. \qquad (3.14b) \end{aligned}$$

Proof. By using the birational canonical transformations l and $\widetilde{l}$, we have

$$\begin{aligned} l(H_i) &= H_i - \frac{q_i p_i}{s_i}, & (3.15)\\ \widetilde{l}(H_i) &= H_i - \frac{1}{s_i}\left(q_i p_i - \frac{\kappa_1 q_i}{g_1 - 1}\right) + \frac{\theta_i}{s_i - 1}, & (3.16)\end{aligned}$$

where $g_1 = \sum_j q_j$. We then obtain the relation between τ-functions and canonical variables:

$$q_i = \frac{s_i \dfrac{\partial}{\partial s_i} \log \dfrac{\widetilde{l}(\tau)}{l(\tau)} - \dfrac{\theta_i s_i}{s_i - 1}}{\displaystyle\sum_j \left(s_j \frac{\partial}{\partial s_j} \log \frac{\widetilde{l}(\tau)}{l(\tau)} - \frac{\theta_j s_j}{s_j - 1}\right) - \kappa_1}, \tag{3.17a}$$

$$q_i\, p_i = s_i \frac{\partial}{\partial s_i} \log \frac{\tau}{l(\tau)}. \tag{3.17b}$$

Here recall the definition of τ-function, $\partial/\partial s_i \log \tau = H_i$. Substitute (3.10) into (3.17) with $s_i = t_i^2$, we get (3.14). ■

3.3 Universal characters

To investigate the special polynomial $T_{m,n}$ in detail, we have to recall the definition of the universal characters; see [7, 16]. For each pair of partitions $[\lambda, \mu] = [(\lambda_1, \lambda_2, \ldots, \lambda_l), (\mu_1, \mu_2, \ldots, \mu_{l'})]$, the *universal character* $S_{[\lambda,\mu]}(x, y)$ is a polynomial in $(x, y) = (x_1, x_2, \ldots, y_1, y_2, \ldots)$ defined as follows:

$$S_{[\lambda,\mu]}(x,y) = \det \begin{pmatrix} q_{\mu_{l'-i+1}+i-j}(y), & 1 \le i \le l' \\ p_{\lambda_{i-l'}-i+j}(x), & l'+1 \le i \le l+l' \end{pmatrix}_{1 \le i,j \le l+l'}. \tag{3.18}$$

Here $p_n(x)$ is determined by the generating function:

$$\sum_{n=0}^{\infty} p_n(x) z^n = e^{\xi(x,z)}, \quad \xi(x,z) = \sum_{n=1}^{\infty} x_n z^n, \tag{3.19}$$

and set $p_{-n}(x) = 0$ for $n > 0$; $q_n(y)$ is the same as $p_n(x)$ except replacing x with y. Note that $p_n(x)$ is explicitly written as follows:

$$p_n(x) = \sum_{k_1 + 2k_2 + \cdots + nk_n = n} \frac{x_1^{k_1} x_2^{k_2} \cdots x_n^{k_n}}{k_1! k_2! \cdots k_n!}. \tag{3.20}$$

If we count the degree of each variable x_n and y_n $(n = 1, 2, \ldots)$ as

$$\deg x_n = n \quad \text{and} \quad \deg y_n = -n,$$

then the universal character $S_{[\lambda,\mu]}(x,y)$ is a weighted homogeneous polynomial of degree $|\lambda|-|\mu|$, where we let $|\lambda|=\lambda_1+\cdots+\lambda_l$. Note that the Schur polynomial $S_\lambda(x)$ (see *e.g.* [8]) is regarded as a special case of the universal character:

$$S_\lambda(x)=\det\left(p_{\lambda_i-i+j}(x)\right)=S_{[\lambda,\emptyset]}(x,y).$$

Example 3.4. When $\lambda=(2,1)$, $\mu=(1)$, the universal character is given as follows:

$$S_{[(2,1),(1)]}(x,y)=\begin{vmatrix} q_1 & q_0 & q_{-1} \\ p_1 & p_2 & p_3 \\ p_{-1} & p_0 & p_1 \end{vmatrix}=y_1\left(\frac{x_1^3}{3}-x_3\right)-x_1^2,$$

which is a weighted homogeneous polynomial of degree $|\lambda|-|\mu|=2$.

The special polynomial $T_{m,n}(t)$ can be written in terms of the universal character.

Theorem 3.5. *The special polynomial $T_{m,n}(t)$ $(m,n\in\mathbb{Z})$ is expressed as follows*:

$$T_{m,n}(t)=N_{m,n}S_{[\lambda,\mu]}(x,y). \tag{3.21}$$

Here $\lambda=(u,u-1,\dots,2,1)$, $\mu=(v,v-1,\dots,2,1)$ with $u=|n-m-1/2|-1/2$, $v=|n+m-1/2|-1/2$; and

$$x_n=\frac{-\kappa_\infty+\sum_i\theta_i t_i^n}{n},\quad y_n=\frac{-\kappa_\infty+\sum_i\theta_i t_i^{-n}}{n}. \tag{3.22}$$

The normalization factor $N_{m,n}$ is given by

$$N_{m,n}=(-1)^{v(v+1)/2}\prod_{i=1}^{N}t_i^{v(v+1)/2}\prod_{j=1}^{u}(2j-1)!!\prod_{k=1}^{v}(2k-1)!!. \tag{3.23}$$

Consequently we have the

Corollary 3.6. *The special polynomial $T_{m,n}(t)$ is indeed a polynomial of degree $m^2+n(n-1)$; furthermore $T_{m,n}(t)\in\mathbb{Z}[\kappa_\infty,\theta_1,\dots,\theta_N][t]$.*

The proof of Theorem 3.5 is given in Sect. 4.

We show in Figure 1 below how the special polynomials $T_{m,n}(t)$ are arranged on (m,n)-lattice. We also give some examples of $T_{m,n}(t)$ of small degrees in the case $N=1$.



Figure 1 Special polynomials $T_{m,n}(t)$.

The special polynomials $T_{m,n}(t)$ for $N=1$ are as follows:

$$\begin{aligned}
&T_{0,0}=T_{0,1}=1,\quad T_{1,0}=T_{-1,1}=-\kappa_\infty+\theta t,\quad T_{1,1}=T_{-1,0}=-\theta+\kappa_\infty t,\\
&T_{0,2}=T_{0,-1}=\kappa_\infty\theta+t-\kappa_\infty^2 t-\theta^2 t+\kappa_\infty\theta t^2,\\
&T_{1,-1}=T_{-1,2}=\kappa_\infty-\kappa_\infty^3+3\kappa_\infty^2\theta t-3\kappa_\infty\theta^2 t^2-\theta t^3+\theta^3 t^3,\\
&T_{1,2}=T_{-1,-1}=\theta-\theta^3+3\kappa_\infty\theta^2 t-3\kappa_\infty^2\theta t^2-\kappa_\infty t^3+\kappa_\infty^3 t^3,\\
&T_{2,0}=T_{-2,1}=-\kappa_\infty\theta+\kappa_\infty^3\theta+4\kappa_\infty^2 t-\kappa_\infty^4 t-3\kappa_\infty^2\theta^2 t-6\kappa_\infty\theta t^2+3\kappa_\infty^3\theta t^2+3\kappa_\infty\theta^3 t^2\\
&\qquad\qquad\qquad +4\theta^2 t^3-3\kappa_\infty^2\theta^2 t^3-\theta^4 t^3-\kappa_\infty\theta t^4+\kappa_\infty\theta^3 t^4.
\end{aligned}$$

Remark 3.7. Under the specialization (3.22), we let $p_n(x)=P_n(t)$. Then the generating function (3.19) is rewritten as follows:

$$\sum_{n=0}^{\infty}P_n(t)z^n=(1-z)^{\kappa_\infty}\prod_i(1-t_i z)^{-\theta_i}. \tag{3.24}$$

Hence $P_n(t)$ has the following expression:

$$P_n(t) = \frac{(-\kappa_\infty)_n}{(1)_n} F_D(-n, \theta_1, \ldots, \theta_N, \kappa_\infty - n + 1; t), \tag{3.25}$$

where F_D denotes the Lauricella hypergeometric series and $(a)_n = a(a+1)(a+2)\cdots(a+n-1)$; see *e.g.* [2, 12, 15].

Remark 3.8. If $N = 1$, $T_{m,n}(t)$ is equivalent to the *Umemura polynomial* of P_{VI}, for which Masuda considered its explicit formula in terms of universal characters; see [10, 11]. We refer also to the results [9] and [17], where a class of rational solutions of P_{V} and that of the (higher order) Painlevé equation of type $A^{(1)}_{2g+1}$ $(g \geq 1)$ are obtained in terms of universal characters.

Remark 3.9. Several other classes of solutions of the Garnier system have been studied. In [15], a family of rational solutions was obtained by the use of Schur polynomials. In [6], solutions in terms of hyperelliptic theta functions were considered from the viewpoint of algebraic geometry.

4 Proof of Theorem 3.5

4.1 A generalization of Jacobi's identity

First we prepare an identity for determinants, which is regarded as a generalization of Jacobi's identity. Let $A = (a_{ij})_{i,j}$ be an $n \times n$ matrix and $\xi^I_J = \xi^I_J(A)$ its minor determinant with respect to rows $I = \{i_1, \ldots, i_r\}$ and columns $J = \{j_1, \ldots, j_r\}$. For two disjoint sets $I, J \subset \{1, \ldots, n\}$, we define $\epsilon(I; J)$ by

$$\epsilon(I; J) = (-1)^{l(I;J)}, \quad l(I; J) = \#\left\{(i, j) \in I \times J \,|\, i > j\right\}. \tag{4.1}$$

Theorem 4.1. *Let* $I = \{1, 2, \ldots, n\}$ *and* $A = (a_{ij})_{i,j \in I}$. *The following quadratic relation among minor determinants of* A *holds*:

$$\xi^I_I \xi^{I-J_1-J_2}_{I-J_1-J_2} = \sum_{\substack{K_1, K_2 \subset I; \\ K_1 \cap (I-J_1-J_2) = \emptyset; \\ K_2 \cap (I-J_1-J_2) = \emptyset}} \epsilon(K_1; K_2) \xi^{I-K_1}_{I-J_1} \xi^{I-K_2}_{I-J_2}, \tag{4.2}$$

where $|J_1| = |K_1| = r_1$ *and* $|J_2| = |K_2| = r_2$.

Let $r_1 = r_2 = 1$, $J_1 = \{1\}$ and $J_2 = \{n\}$, then (4.2) recovers Jacobi's identity (see [3]):

$$\xi^{1\cdots n}_{1\cdots n} \xi^{2\cdots n-1}_{2\cdots n-1} = \xi^{2\cdots n}_{2\cdots n} \xi^{1\cdots n-1}_{1\cdots n-1} - \xi^{1\cdots n-1}_{2\cdots n} \xi^{2\cdots n}_{1\cdots n-1}, \tag{4.3}$$

in fact.

Proof of Theorem 4.1. Without loss of generality, we can set $J_1 = \{1, 2, \ldots, r_1\}$ and $J_2 = \{n-r_2+1, \ldots, n-1, n\}$. Let $\widetilde{I} = \{1, 2, \ldots, 2n-r_1-r_2\}$. Consider a $(2n-r_1-r_2)\times(2n-r_1-r_2)$ matrix $B = (b_{ij})_{i,j\in\widetilde{I}}$ given as follows:

$$\begin{array}{llll} \text{(i)} & b_{ij} = a_{ij} & \text{for} & i, j \in I; \\ \text{(ii)} & b_{ij} = a_{i,j-n+r_1} & \text{for} & i \in I,\ j \in \widetilde{I} \setminus I; \\ \text{(iii)} & b_{ij} = a_{i-n+r_1,j} & \text{for} & i \in \widetilde{I} \setminus I,\ j \in J_1; \\ \text{(iv)} & b_{ij} = 0 & \text{for} & i \in \widetilde{I} \setminus I,\ j \in I \setminus J_1; \\ \text{(v)} & b_{ij} = a_{i-n+r_1,j-n+r_1} & \text{for} & i \in \widetilde{I} \setminus I,\ j \in \widetilde{I} \setminus I, \end{array} \tag{4.4}$$

i.e., write A as

$$A = \left[\begin{array}{c|c|c} A_{11} & A_{12} & A_{13} \\ \hline A_{21} & \boldsymbol{A}_{22} & A_{23} \\ \hline A_{31} & A_{32} & A_{33} \end{array}\right],$$

then B is written as

$$B = \left[\begin{array}{c|c|c|c} A_{11} & A_{12} & A_{13} & A_{12} \\ \hline A_{21} & \boldsymbol{A}_{22} & A_{23} & \boldsymbol{A}_{22} \\ \hline A_{31} & A_{32} & A_{33} & A_{32} \\ \hline A_{21} & 0 & 0 & \boldsymbol{A}_{22} \end{array}\right].$$

Apply the Laplace expansion with respect to rows I and rows $\widetilde{I} \setminus I$, we obtain

$$\det B = \xi^{I}_{I} \xi^{I-J_1-J_2}_{I-J_1-J_2}. \tag{4.5}$$

On the other hand, by the Laplace expansion with respect to columns $I \setminus J_1$ and columns $(\widetilde{I} \setminus I) \cup J_1$, we have

$$\det B = \sum_{\substack{K_1, K_2 \subset I; \\ K_1 \cap (I-J_1-J_2)=\emptyset; \\ K_2 \cap (I-J_1-J_2)=\emptyset}} \epsilon(K_1; K_2) \xi^{I-K_1}_{I-J_1} \xi^{I-K_2}_{I-J_2}. \tag{4.6}$$

Thus we verify (4.2). ■

4.2 Vertex operators

Introduce the vertex operators $V_m(k; x, y)$ $(m \in \mathbb{Z})$ defined by (see [16])

$$V_m(k; x, y) = e^{m\xi(x-\widetilde{\partial}_y, k)} e^{-m\xi(\widetilde{\partial}_x, k^{-1})}, \tag{4.7}$$

where $\widetilde{\partial}_x$ stands for $\left(\frac{\partial}{\partial x_1}, \frac{1}{2}\frac{\partial}{\partial x_2}, \frac{1}{3}\frac{\partial}{\partial x_3}, \ldots\right)$ and $\xi(x, k) = \sum_{n=1}^{\infty} x_n k^n$. Define the differential operators X_n and Y_n $(n \in \mathbb{Z})$ by

$$\begin{aligned} X(k) &= \textstyle\sum_{n\in\mathbb{Z}} X_n k^n = V_1(k; x, y), \\ Y(k) &= \textstyle\sum_{n\in\mathbb{Z}} Y_n k^{-n} = V_1(k^{-1}; y, x). \end{aligned} \tag{4.8}$$

We have the following lemmas; see [16].

Lemma 4.2. *The operators X_n and Y_n $(n \in \mathbb{Z})$ are raising operators for the universal characters in the sense that*

$$S_{[\lambda,\mu]}(x,y) = X_{\lambda_1} \cdots X_{\lambda_l} Y_{\mu_1} \cdots Y_{\mu_{l'}} \cdot 1. \tag{4.9}$$

Lemma 4.3. *The following relations hold:*

$$\begin{aligned} &X_m X_n + X_{n-1} X_{m+1} = 0, \\ &Y_m Y_n + Y_{n-1} Y_{m+1} = 0, \\ &[X_m, Y_n] = 0, \end{aligned} \tag{4.10}$$

for $m, n \in \mathbb{Z}$. In particular $X_n X_{n+1} = Y_n Y_{n+1} = 0$.

4.3 Proof of Theorem 3.5

Introduce the Euler operator

$$E = \sum_{n=1}^{\infty} \left(n x_n \frac{\partial}{\partial x_n} - n y_n \frac{\partial}{\partial y_n} \right), \tag{4.11}$$

and operators L^+, L^- given as follows:

$$L^+ = \frac{x_1^2}{2} + \sum_{n=1}^{\infty} \left((n+2) x_{n+2} \frac{\partial}{\partial x_n} - n y_n \frac{\partial}{\partial y_{n+2}} \right) - x_1 \frac{\partial}{\partial y_1} - \left(-\kappa_\infty + \sum_i \theta_i \right) \frac{\partial}{\partial y_2}, \tag{4.12}$$

$$L^- = \frac{y_1^2}{2} + \sum_{n=1}^{\infty} \left((n+2) y_{n+2} \frac{\partial}{\partial y_n} - n x_n \frac{\partial}{\partial x_{n+2}} \right) - y_1 \frac{\partial}{\partial x_1} - \left(-\kappa_\infty + \sum_i \theta_i \right) \frac{\partial}{\partial x_2}. \tag{4.13}$$

Note that E, L^+, and L^- are homogeneous operators of degrees 0, 2, and -2, respectively. Consider the change of the variables

$$x_n = \frac{-\kappa_\infty + \sum_i \theta_i t_i^n}{n}, \quad y_n = \frac{-\kappa_\infty + \sum_i \theta_i t_i^{-n}}{n}, \tag{4.14}$$

and

$$\widetilde{T}_{m,n}(x,y) = (-1)^{-v(v+1)/2} \prod_i t_i^{-v(v+1)/2} T_{m,n}(t), \tag{4.15}$$

where $u = |n-m-1/2| - 1/2$, $v = |n+m-1/2| - 1/2$. Substitute this into (3.11), we have the recurrence relations for $\widetilde{T}_{m,n}(x,y)$:

$$\begin{aligned} &-\widetilde{T}_{m+1,n}\, \widetilde{T}_{m-1,n} \\ &\quad = \left\{ \left(L^- + E - \frac{y_1^2}{2} - 2 \right) \left(L^+ - E - \frac{x_1^2}{2} \right) \log \widetilde{T}_{m,n} - x_1 y_1 + (2m)^2 \right\} {\widetilde{T}_{m,n}}^2, \end{aligned} \tag{4.16a}$$

$$\begin{aligned} &-\widetilde{T}_{m,n+1}\, \widetilde{T}_{m,n-1} \\ &\quad = \left\{ \left(L^- + E - \frac{y_1^2}{2} - 2 \right) \left(L^+ - E - \frac{x_1^2}{2} \right) \log \widetilde{T}_{m,n} - x_1 y_1 + (2n-1)^2 \right\} {\widetilde{T}_{m,n}}^2, \end{aligned} \tag{4.16b}$$

where the initial values are given by

$$\widetilde{T}_{0,0} = \widetilde{T}_{0,1} = 1, \quad \widetilde{T}_{1,0} = x_1, \quad \widetilde{T}_{1,1} = y_1. \tag{4.17}$$

Note that we have

$$\widetilde{T}_{-m,n}(x,y) = \widetilde{T}_{m,1-n}(x,y) = \widetilde{T}_{m,n}(y,x), \tag{4.18}$$

from (3.13).

Theorem 3.5 follows immediately from the

Proposition 4.4. *Let*

$$\widetilde{T}_{m,n}(x,y) = \prod_{j=1}^{u}(2j-1)!! \prod_{k=1}^{v}(2k-1)!!\, S_{[\lambda,\mu]}(x,y), \tag{4.19}$$

where $\lambda = (u, u-1, \dots, 2, 1)$ *and* $\mu = (v, v-1, \dots, 2, 1)$*, then* $\widetilde{T}_{m,n}(x,y)$ *satisfies* (4.16) *and* (4.17)*.*

We prepare some lemmas to verify Proposition 4.4.

Lemma 4.5. *The following commutation relations hold for* $n \in \mathbb{Z}$*:*

$$[X_n, L^+] = -\left(n + \frac{3}{2}\right) X_{n+2} + 2\left(x_2 - \frac{\partial}{\partial y_2}\right) X_n, \tag{4.20}$$

$$[Y_n, L^+] = \left(n - \frac{3}{2} - \kappa_\infty + \sum_i \theta_i\right) Y_{n-2} - Y_n \frac{\partial}{\partial y_2}, \tag{4.21}$$

$$[X_n, x_2] = -\frac{1}{2} X_{n+2}, \tag{4.22}$$

$$[Y_n, x_2] = -\frac{1}{2} Y_{n-2}. \tag{4.23}$$

Proof. Notice that for any operators A and B,

$$e^A B e^{-A} = e^{\mathrm{ad}(A)} B = B + [A,B] + \frac{1}{2!}[A,[A,B]] + \cdots,$$

where $\mathrm{ad}(A)(B) = [A,B]$. We have

$$[\xi(x - \widetilde{\partial}_y, k), L^+] = -\sum_{m=1}^{\infty} \left\{ (m+2)x_{m+2} - \frac{\partial}{\partial y_{m+2}} \right\} k^m,$$

so that

$$[e^{\xi(x-\widetilde{\partial}_y,k)}, L^+] = -\sum_{m=1}^{\infty} \left\{ (m+2)x_{m+2} - \frac{\partial}{\partial y_{m+2}} \right\} k^m e^{\xi(x-\widetilde{\partial}_y,k)}. \tag{4.24}$$

On the other hand, we have

$$\begin{aligned}[-\xi(\widetilde{\partial}_x,k^{-1}),L^+] &= -\left(x_1-\frac{\partial}{\partial y_1}\right)k^{-1}-\sum_{m=1}^{\infty}k^{-m-2}\frac{\partial}{\partial x_m},\\ [-\xi(\widetilde{\partial}_x,k^{-1}),[-\xi(\widetilde{\partial}_x,k^{-1}),L^+]] &= k^{-2},\end{aligned}$$

then

$$[e^{-\xi(\widetilde{\partial}_x,k^{-1})},L^+]=\left\{-\left(x_1-\frac{\partial}{\partial y_1}\right)k^{-1}+\frac{k^{-2}}{2}-\sum_{m=1}^{\infty}k^{-m-2}\frac{\partial}{\partial x_m}\right\}e^{-\xi(\widetilde{\partial}_x,k^{-1})}. \tag{4.25}$$

Noticing

$$k^{-1}\frac{\partial}{\partial k}X(k)=\sum_{m=1}^{\infty}\left(mx_m-\frac{\partial}{\partial y_m}\right)k^{m-2}X(k)+e^{\xi(x-\widetilde{\partial}_y,k)}\sum_{m=1}^{\infty}k^{-m-2}\frac{\partial}{\partial x_m}e^{-\xi(\widetilde{\partial}_x,k^{-1})},$$

from (4.24) and (4.25), we obtain

$$\begin{aligned}[X(k),L^+] &= e^{\xi(x-\widetilde{\partial}_y,k)}[e^{-\xi(\widetilde{\partial}_x,k^{-1})},L^+]+[e^{\xi(x-\widetilde{\partial}_y,k)},L^+]e^{-\xi(\widetilde{\partial}_x,k^{-1})}\\ &= \left\{-k^{-1}\frac{\partial}{\partial k}+\frac{k^{-2}}{2}+2\left(x_2-\frac{1}{2}\frac{\partial}{\partial y_2}\right)\right\}X(k). \end{aligned}\tag{4.26}$$

Take the coefficient of k^n, we verify (4.20).

We have

$$\begin{aligned}&[\xi(y-\widetilde{\partial}_x,k^{-1}),L^+]=k^{-1}\frac{\partial}{\partial y_1}+\left(-\kappa_\infty+\sum_i\theta_i\right)k^{-2}+\sum_{m=1}^{\infty}\left(my_m-\frac{\partial}{\partial x_m}\right)k^{-m-2},\\ &[\xi(y-\widetilde{\partial}_x,k^{-1}),[\xi(y-\widetilde{\partial}_x,k^{-1}),L^+]]=-k^{-2},\\ &[-\xi(\widetilde{\partial}_y,k),L^+]=\sum_{m=1}^{\infty}k^m\frac{\partial}{\partial y_{m+2}},\end{aligned}$$

so that

$$\begin{aligned}[e^{\xi(y-\widetilde{\partial}_x,k^{-1})},L^+] &= \left\{k^{-1}\frac{\partial}{\partial y_1}+\left(-\kappa_\infty+\sum_i\theta_i-\frac{1}{2}\right)k^{-2}+\sum_{m=1}^{\infty}\left(my_m-\frac{\partial}{\partial x_m}\right)k^{-m-2}\right\},\\ [e^{-\xi(\widetilde{\partial}_y,k)},L^+] &= \sum_{m=1}^{\infty}k^m\frac{\partial}{\partial y_{m+2}}e^{-\xi(\widetilde{\partial}_y,k)}.\end{aligned}$$

Thus we obtain

$$\begin{aligned}[Y(k),L^+] &= e^{\xi(y-\widetilde{\partial}_x,k^{-1})}[e^{-\xi(\widetilde{\partial}_y,k)},L^+]+[e^{\xi(y-\widetilde{\partial}_x,k^{-1})},L^+]e^{-\xi(\widetilde{\partial}_y,k)}\\ &= \left\{-k^{-1}\frac{\partial}{\partial k}+\left(-\kappa_\infty+\sum_i\theta_i+\frac{1}{2}\right)k^{-2}\right\}Y(k)-Y(k)\frac{\partial}{\partial y_2}, \end{aligned}\tag{4.27}$$

whose coefficient of k^{-n} yields (4.21).

By $[-\xi(\widetilde{\partial}_x, k^{-1}), x_2] = -k^{-2}/2$, we have

$$[e^{-\xi(\widetilde{\partial}_x, k^{-1})}, x_2] = -\frac{k^{-2}}{2} e^{-\xi(\widetilde{\partial}_x, k^{-1})},$$

therefore

$$[X(k), x_2] = -\frac{k^{-2}}{2} X(k), \quad [Y(k), x_2] = -\frac{k^{-2}}{2} Y(k). \tag{4.28}$$

Take the coefficients of k^n and k^{-n}, we obtain (4.22) and (4.23) respectively. ■

Lemma 4.6. *For integers $u, v \geq 0$, the following formulae hold:*

$$L^+ S_{[u!,v!]}(x,y) = (2u+1) S_{[(u+2,u-1,\ldots,1),v!]}(x,y) - (2u+1) x_2 S_{[u!,v!]}(x,y), \tag{4.29}$$

$$L^- S_{[u!,v!]}(x,y) = (2v+1) S_{[u!,(v+2,v-1,\ldots,1)]}(x,y) - (2v+1) y_2 S_{[u!,v!]}(x,y), \tag{4.30}$$

$$\begin{aligned} L^+ S_{[u!,(v+2,v-1,\ldots,1)]}(x,y) = {} & (2u+1) S_{[(u+2,u-1,\ldots,1),(v+2,v-1,\ldots,1)]}(x,y) \\ & -(2u+1) x_2 S_{[u!,(v+2,v-1,\ldots,1)]}(x,y) \\ & - \left(v - u - \kappa_\infty + \sum_i \theta_i \right) S_{[u!,v!]}(x,y), \end{aligned} \tag{4.31}$$

$$\begin{aligned} L^- S_{[(u+2,u-1,\ldots,1),v!]}(x,y) = {} & (2v+1) S_{[(u+2,u-1,\ldots,1),(v+2,v-1,\ldots,1)]}(x,y) \\ & -(2v+1) y_2 S_{[(u+2,u-1,\ldots,1),v!]}(x,y) \\ & - \left(u - v - \kappa_\infty + \sum_i \theta_i \right) S_{[u!,v!]}(x,y). \end{aligned} \tag{4.32}$$

Here $u! = (u, u-1, \ldots, 2, 1)$.

Proof. First we shall show that

$$L^+ S_{[u!,\emptyset]}(x,y) = (2u+1) S_{[(u+2,u-1,\ldots,1),\emptyset]}(x,y) - (2u+1) x_2 S_{[u!,\emptyset]}(x,y), \tag{4.33}$$

by induction. Using $S_{[\emptyset,\emptyset]}(x,y) = 1$ and $S_{[(2),\emptyset]}(x,y) = x_1^2/2 + x_2$, it is easy to verify for $u = 0$. Assume that (4.33) is true for $u-1$. Applying X_u, we have

$$\begin{aligned} X_u L^+ S_{[(u-1)!,\emptyset]}(x,y) &= L^+ S_{[u!,\emptyset]}(x,y) + [X_u, L^+] S_{[(u-1)!,\emptyset]}(x,y) \\ &= (L^+ + 2x_2) S_{[u!,\emptyset]}(x,y) - \left(u + \frac{3}{2} \right) S_{[(u+2,u-1,\ldots,1),\emptyset]}(x,y), \end{aligned}$$

and

$$\begin{aligned} & X_u \left((2u-1) S_{[(u+1,u-2,\ldots,1),\emptyset]}(x,y) - (2u-1) x_2 S_{[(u-1)!,\emptyset]}(x,y) \right) \\ & \quad = -(2u-1) x_2 S_{[u!,\emptyset]}(x,y) + \frac{1}{2}(2u-1) S_{[u!,\emptyset]}(x,y), \end{aligned}$$

by using the commutation relations (4.20) and the property $X_kX_{k+1}=0$. Then, by the assumption, we have the desired equation (4.33) immediately. Applying $Y_vY_{v-1}\cdots Y_1$ to (4.33) we obtain (4.29). Here we recall the commutation relations (4.21), (4.23), and $Y_kY_{k+1}=0$.

Since L^- is the same as L^+ except exchanging x with y, we verify (4.30) immediately.

Notice that $S_{[u!,v!]}(x,y)$ does not depend on y_{2n} $(n=1,2,\ldots)$. Applying Y_{v+3} to (4.29), we have

$$Y_{v+3}L^+S_{[u!,v!]}(x,y)=L^+S_{[u!,(v+3,v,\ldots,1)]}(x,y)+\left(v+\frac{3}{2}-\kappa_\infty+\sum_i\theta_i\right)S_{[u!,(v+1)!]}(x,y),$$

and

$$\begin{aligned}&Y_{v+3}\left((2u+1)S_{[(u+2,u-1,\ldots,1),v!]}(x,y)-(2u+1)x_2S_{[u!,v!]}(x,y)\right)\\&\quad=(2u+1)S_{[(u+2,u-1,\ldots,1),(v+3,v,\ldots,1)]}(x,y)-(2u+1)x_2S_{[u!,(v+3,v,\ldots,1)]}(x,y)\\&\qquad+\left(u+\frac{1}{2}\right)S_{[u!,(v+1)!]}(x,y).\end{aligned}$$

Thus we verify (4.31). Similarly (4.32) also holds. ■

Proof of Proposition 4.4. For the sake of simplicity, we use the following notations:

$$\begin{aligned}S&=S_{[u!,v!]}(x,y),\\S^+&=S_{[(u+2,u-1,\ldots,1),v!]}(x,y),\\S^-&=S_{[u!,(v+2,v-1,\ldots,1)]}(x,y),\\S^{+-}&=S_{[(u+2,u-1,\ldots,1),(v+2,v-1,\ldots,1)]}(x,y).\end{aligned}\tag{4.34}$$

We have

$$\begin{aligned}&\left(\left(L^-+E-\frac{y_1^2}{2}-2\right)\left(L^+-E-\frac{x_1^2}{2}\right)\log S\right)S^2\\&\qquad=\left(L^-+E-\frac{y_1^2}{2}\right)\left(L^+-E-\frac{x_1^2}{2}\right)S\cdot S\\&\qquad\quad-\left(L^-+E-\frac{y_1^2}{2}\right)S\cdot\left(L^+-E-\frac{x_1^2}{2}\right)S\\&\qquad\quad-2\left(L^+-E-\frac{x_1^2}{2}\right)S\cdot S.\end{aligned}\tag{4.35}$$

Since $S_{[\lambda,\mu]}(x,y)$ is a weighted homogeneous polynomial of degree $|\lambda|-|\mu|$, the Euler operator E acts on it as

$$ES_{[\lambda,\mu]}(x,y)=(|\lambda|-|\mu|)S_{[\lambda,\mu]}(x,y).\tag{4.36}$$

Then by Lemma 4.6 we have

$$\begin{aligned}&\left(\left(L^-+E-\frac{y_1^2}{2}-2\right)\left(L^+-E-\frac{x_1^2}{2}\right)\log S-x_1y_1\right)S^2\\&\qquad=(2u+1)(2v+1)S^{+-}S-(2u+1)(2v+1)S^+S^--(u-v)^2S^2.\end{aligned}\tag{4.37}$$

Now let us substitute (4.19) into the recurrence relations (4.16). By virtue of (4.18), it is enough to consider the cases (I) $n-m-1/2>0$, $n+m-1/2>0$; and (II) $n-m-1/2<0$, $n+m-1/2>0$.

First we deal with the case (I), that is, $m=(v-u)/2$, $n=(u+v+2)/2$. Substitute (4.19) into the both sides of (4.16), we have

$$\begin{aligned}\text{LHS of (4.16a)} &= -(2u+1)(2v+1)C_{u,v}S_{[(u+1)!,(v-1)!]}\cdot S_{[(u-1)!,(v+1)!]},\\ \text{RHS of (4.16a)} &= (2u+1)(2v+1)C_{u,v}(S^{+-}S-S^{+}S^{-}),\end{aligned}$$

and

$$\begin{aligned}\text{LHS of (4.16b)} &= -(2u+1)(2v+1)C_{u,v}S_{[(u+1)!,(v+1)!]}\cdot S_{[(u-1)!,(v-1)!]},\\ \text{RHS of (4.16b)} &= (2u+1)(2v+1)C_{u,v}(S^{+-}S-S^{+}S^{-}+S^{2}),\end{aligned}$$

respectively. Here we put $C_{u,v}=\left(\prod_{j=1}^{u}(2j-1)!!\prod_{k=1}^{v}(2k-1)!!\right)^2$. Thus it is sufficient to prove

$$\begin{aligned}-S_{[(u+1)!,(v-1)!]}\cdot S_{[(u-1)!,(v+1)!]} &= S^{+-}S-S^{+}S^{-}, && (4.38)\\ -S_{[(u+1)!,(v+1)!]}\cdot S_{[(u-1)!,(v-1)!]} &= S^{+-}S-S^{+}S^{-}+S^{2}. && (4.39)\end{aligned}$$

By using Lemma 4.7 below, we immediately verify (4.38) and (4.39).

The verification for the case (II) is the same. ■

Lemma 4.7. *The following formulae hold*:

$$S_{[(u+1)!,(v+1)!]}\cdot S_{[(u-1)!,(v-1)!]}-S_{[(u+1)!,(v-1)!]}\cdot S_{[(u-1)!,(v+1)!]}+{S_{[u!,v!]}}^{2}=0, \tag{4.40}$$

$$\begin{aligned}S_{[(u+1)!,(v-1)!]}\cdot S_{[(u-1)!,(v+1)!]}-S_{[u!,(v+2,v-1,\ldots,1)]}\cdot S_{[(u+2,u-1,\ldots,1),v!]}&\\ +S_{[(u+2,u-1,\ldots,1),(v+2,v-1,\ldots,1)]}\cdot S_{[u!,v!]}=0.&\end{aligned} \tag{4.41}$$

Proof. Consider a $(u+v+2)\times(u+v+2)$ matrix

$$M=\left[\begin{array}{cccccccccccc}
q_1 & q_0 & 0 & 0 & \cdots & & & & \cdots & 0 & 0 & 0\\ \cline{3-10}
 & q_2 & \multicolumn{1}{|c}{q_1} & & & & & & & \multicolumn{1}{c|}{} & & \\
 & & \multicolumn{1}{|c}{q_3} & q_2 & & & & & & \multicolumn{1}{c|}{} & & \\
 & & \multicolumn{1}{|c}{} & \ddots & \ddots & & & & & \multicolumn{1}{c|}{} & & \\
 & & \multicolumn{1}{|c}{} & & q_v & q_{v-1} & & & & \multicolumn{1}{c|}{} & & \\ \cline{3-10}
 & & & & \cdots & q_{v+1} & q_v & \cdots & & & & \\
 & & & & \cdots & p_u & p_{u+1} & \cdots & & & & \\ \cline{3-10}
 & & \multicolumn{1}{|c}{} & & & & p_{u-1} & p_u & & \multicolumn{1}{c|}{} & & \\
 & & \multicolumn{1}{|c}{} & & & & & \ddots & \ddots & \multicolumn{1}{c|}{} & & \\
 & & \multicolumn{1}{|c}{} & & & & & & p_2 & \multicolumn{1}{c|}{p_3} & & \\
 & & \multicolumn{1}{|c}{} & & & & & & & \multicolumn{1}{c|}{p_1} & p_2 & \\ \cline{3-10}
0 & 0 & 0 & \cdots & & & & \cdots & 0 & 0 & p_0 & p_1
\end{array}\right], \tag{4.42}$$

so that $D = \det M = S_{[(u+1)!,(v+1)!]}(x,y)$. Denote by $D[i_1, i_2, \ldots; j_1, j_2, \ldots]$ its minor determinant removing rows $\{i_k\}$ and columns $\{j_k\}$. It is easy to see that

$$\begin{aligned}
&D\,[1, v+1, v+2, u+v+2; 1, 2, u+v+1, u+v+2] = S_{[(u-1)!,(v-1)!]}(x,y),\\
&D[1, v+1; 1, 2] = S_{[(u+1)!,(v-1)!]}(x,y),\\
&D[v+2, u+v+2; u+v+1, u+v+2] = S_{[(u-1)!,(v+1)!]}(x,y),\\
&D[1, v+2; 1, 2] = D[v+1, u+v+2; u+v+1, u+v+2] = S_{[u!,v!]}(x,y).
\end{aligned} \tag{4.43}$$

Applying Theorem 4.1, we have

$$\begin{aligned}
&DD[1, v+1, v+2, u+v+2; 1, 2, u+v+1, u+v+2]\\
&\quad = D[1, v+1; 1, 2]D[v+2, u+v+2; u+v+1, u+v+2]\\
&\qquad - D[1, v+2; 1, 2]D[v+1, u+v+2; u+v+1, u+v+2],
\end{aligned} \tag{4.44}$$

which coincides with (4.40).

Take a $(u+v+2) \times (u+v)$ matrix

$$\widetilde{M} = \left[\begin{array}{cccccccc}
\hline
q_1 & q_0 & 0 & \cdots & & & \cdots & 0\\
 & \ddots & \ddots & & & & & \\
 & & q_{v-1} & q_{v-2} & & & & \\
\hline
 & & \cdots & q_v & q_{v-1} & \cdots & & \\
 & & \cdots & q_{v+2} & q_{v+1} & \cdots & & \\
 & & \cdots & p_{u+1} & p_{u+2} & \cdots & & \\
 & & \cdots & p_{u-1} & p_u & \cdots & & \\
\hline
 & & & & p_{u-2} & p_{u-1} & & \\
 & & & & & \ddots & \ddots & \\
0 & \cdots & & & \cdots & 0 & p_0 & p_1\\
\hline
\end{array}\right], \tag{4.45}$$

then

$$\begin{aligned}
&D[v, v+1; \emptyset] = S_{[(u+1)!,(v-1)!]}(x,y), \quad D[v+2, v+3; \emptyset] = S_{[(u-1)!,(v+1)!]}(x,y),\\
&D[v, v+2; \emptyset] = S_{[u!,(v+2,v-1,\ldots,1)]}(x,y), \quad D[v+1, v+3; \emptyset] = S_{[(u+2,u-1,\ldots,1),v!]}(x,y),\\
&D[v, v+3; \emptyset] = S_{[(u+2,u-1,\ldots,1),(v+2,v-1,\ldots,1)]}(x,y), \quad D[v+1, v+2; \emptyset] = S_{[u!,v!]}(x,y).
\end{aligned} \tag{4.46}$$

By the Plücker relation, we have

$$\begin{aligned}
&D[v, v+1; \emptyset]D[v+2, v+3; \emptyset] - D[v, v+2; \emptyset]D[v+1, v+3; \emptyset]\\
&\qquad + D[v, v+3; \emptyset]D[v+1, v+2; \emptyset] = 0,
\end{aligned} \tag{4.47}$$

which coincides with (4.41). ■

Acknowledgement. The author wishes to thank Professor Kazuo Okamoto for valuable discussions. This work is partially supported by a fellowship of the Japan Society for the Promotion of Science (JSPS).

References

[1] Garnier, R.: Sur des équations différentielles du troisième ordre dont l'intégrale générale est uniforme et sur un classe d'équations nouvelles d'order supérieur dont l'intégrale générale a ses points critiques fixes. Ann. Sci. École Norm. Sup. (3) **29**, 1–126 (1912)

[2] Iwasaki, K., Kimura, H., Shimomura, S., Yoshida, M.: *From Gauss to Painlevé: a modern theory of special functions.* Aspects of Mathematics, vol. E16, Braunschweig: Vieweg Verlag, 1991

[3] Jacobi, C. G. J.: De formatione et proprietatibus determinantium. J. reine angew. Math. **22**, 285–318 (1841)

[4] Kimura, H., Okamoto, K.: On the polynomial Hamiltonian structure of the Garnier system. J. Math. Pures Appl. (9) **63**, 129–146 (1984)

[5] Kimura, H.: Symmetries of the Garnier system and of the associated polynomial Hamiltonian system. Proc. Japan Acad. Ser. A Math. Sci. **66**, 176–178 (1990)

[6] Kitaev, A. V., Korotkin, D. A.: On solutions of the Schlesinger equations in terms of Θ-functions. Internat. Math. Res. Notices **1998** no. 17, 877–905 (1998)

[7] Koike, K.: On the decomposition of tensor products of the representations of the classical groups: By means of the universal characters. Adv. Math. **74**, 57–86 (1989)

[8] Macdonald, I. G.: *Symmetric Functions and Hall Polynomials.* 2nd ed., Oxford Mathematical Monographs, New York: Oxford University Press Inc., 1995

[9] Masuda, T., Ohta, Y., Kajiwara, K.: A determinant formula for a class of rational solutions of Painlevé V equation. Nagoya Math. J. **168**, 1–25 (2002)

[10] Masuda, T.: On a class of algebraic solutions to the Painlevé VI equation, its determinant formula and coalescence cascade. Funkcial. Ekvac. **46**, 121–171 (2003)

[11] Noumi, M., Okada, S., Okamoto, K., Umemura, H.: *Special polynomials associated with the Painlevé equations II.* In: Integrable Systems and Algebraic Geometry, eds. Saito, M.-H., Shimizu, Y., Ueno, K., Singapore: World Scientific, 1998, pp. 349–372

[12] Okamoto K., Kimura, H.: On particular solutions of the Garnier systems and the hypergeometric functions of several variables. Quart. J. Math. Oxford Ser. (2) **37**, 61–80 (1986)

[13] Okamoto, K.: Studies on the Painlevé equations, I. Ann. Mat. Pura Appl. (4) **146**, 337–381 (1987)

[14] Tsuda, T.: Birational symmetries, Hirota bilinear forms and special solutions of the Garnier systems in 2-variables. J. Math. Sci. Univ. Tokyo **10**, 355–371 (2003)

[15] Tsuda, T.: Rational solutions of the Garnier system in terms of Schur polynomials. Internat. Math. Res. Notices **2003** no. 43, 2341–2358 (2003)

[16] Tsuda, T.: Universal characters and an extension of the KP hierarchy. Commun. Math. Phys. **248**, 501–526 (2004)

[17] Tsuda, T.: *Universal characters, integrable chains and the Painlevé equations*. Submitted to Adv. Math., Preprint: UTMS 2004–14

[18] Tsuda, T.: *Universal characters and q-Painlevé systems*. Submitted to Commun. Math. Phys.

[19] Tsuda, T.: *Universal characters and integrable systems*. Ph.D. thesis, The University of Tokyo, 2003